

## DIS 12B

### 1 Sum of Independent Gaussians

In this question, we will introduce an important property of the Gaussian distribution: the sum of independent Gaussians is also a Gaussian.

Let  $X$  and  $Y$  be independent standard Gaussian random variables. Recall that the density of the standard Gaussian is

$$f(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right).$$

- (a) What is the joint density of  $X$  and  $Y$ ?
- (b) Observe that the joint density of  $X$  and  $Y$ ,  $f_{X,Y}(x,y)$ , only depends on the quantity  $x^2 + y^2$ , which is the distance from the origin. In other words, the Gaussian is *rotationally symmetric*. Next, we will try to find the density of  $X + Y$ . To do this, draw a picture of the Cartesian plane and draw the region  $x + y \leq c$ , where  $c$  is a real number of your choice.
- (c) Now, rotate your picture clockwise by  $\pi/4$  so that the line  $X + Y = c$  is now vertical. Redraw your figure. Let  $X'$  and  $Y'$  denote the random variables which correspond to the  $\pi/4$  clockwise rotation of  $(X, Y)$  and express the new shaded region in terms of  $X'$  and  $Y'$ .
- (d) By rotational symmetry of the Gaussian,  $(X', Y')$  has the same distribution as  $(X, Y)$ . Argue that  $X + Y$  has the same distribution as  $\sqrt{2}Z$ , where  $Z$  is a standard Gaussian. This proves the following important fact: *the sum of independent Gaussians is also a Gaussian*. Notice that  $X \sim \mathcal{N}(0, 1)$ ,  $Y \sim \mathcal{N}(0, 1)$  and  $Z \sim \mathcal{N}(0, 2)$ . In general, if  $X$  and  $Y$  are independent Gaussians, then  $X + Y$  is a Gaussian with mean  $\mu_X + \mu_Y$  and variance  $\sigma_X^2 + \sigma_Y^2$ .

(e) Recall the CLT:

If  $\{X_i\}_{i \in \mathbb{N}}$  is a sequence of i.i.d. random variables with mean  $\mu \in \mathbb{R}$  and variance  $\sigma^2 < \infty$ , then:

$$\frac{X_1 + \cdots + X_n - n\mu}{\sigma\sqrt{n}} \xrightarrow{\text{in distribution}} \mathcal{N}(0, 1) \quad \text{as } n \rightarrow \infty.$$

Prove that the CLT holds for the special case when the  $X_i$  are i.i.d.  $\mathcal{N}(0, 1)$ .

## 2 Inequality Practice

(a)  $X$  is a random variable such that  $X > -5$  and  $\mathbb{E}[X] = -3$ . Find an upper bound for the probability of  $X$  being greater than or equal to  $-1$ .

(b) You roll a die 100 times. Let  $Y$  be the sum of the numbers that appear on the die throughout the 100 rolls. Use Chebyshev's inequality to bound the probability of the sum  $Y$  being greater than 400 or less than 300.

## 3 Poisson Confidence Interval

You collect  $n$  samples ( $n$  is a positive integer)  $X_1, \dots, X_n$ , which are i.i.d. and known to be drawn from a Poisson distribution (with unknown mean). However, you have a bound on the mean: from a confidential source, you know that  $\lambda \leq 2$ . Find a  $1 - \delta$  confidence interval ( $\delta \in (0, 1)$ ) for  $\lambda$  using Chebyshev's Inequality.